

KEYWORD SEARCH

GO

OUR PRODUCTS

STORE LOCATOR

ABOUT US

FAQs

LINKS

CONTACT US

**B.F. ASCHER & CO. INC.**

pharmaceuticals

*Makers of Consumer Health Care
Products for Better Living.*



Cough-X Lozenges

- Dual acting
- Pleasant tasting
- Dye free

[LOCATE THIS PRODUCT IN A STORE NEAR YOU](#)
[Click to return to the PRODUCT CATEGORY PAGE](#)

Cough syrup effectiveness in a convenient lozenge form

Cough-X contains dextromethorphan (the same cough suppressant used in most popular cough liquids) and benzocaine (for sore throat pain). Pleasant menthol eucalyptus flavor. Temporarily suppresses cough due to minor throat and bronchial irritants. Also, for the temporary relief of occasional minor irritation and sore throat.

INGREDIENTS

Active Ingredients: Each lozenge contains dextromethorphan hydrobromide 5 mg and benzocaine 2 mg.

Also contains: Corn syrup, eucalyptus oil, menthol, propylene glycol and sucrose.

DIRECTIONS

Allow lozenge to dissolve slowly in the mouth. Adults and children 6 years of age and older: One lozenge every 2 hours as needed not to exceed 12 lozenges in 24 hours or as directed by a physician. Children 2 to 6 years of age: One lozenge every 4 hours not to exceed 6 lozenges in 24 hours or as directed by a physician. Children under 2 years of age: consult a physician.

In children, take care to prevent choking on lozenge.

WARNINGS

A persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur or if sore throat is severe, persists for more than 2 days or if cough and/or sore throat is accompanied or followed by fever, persistent headache, rash, swelling, nausea or vomiting, consult a physician. Do not take this product for persistent or chronic cough such as occurs with smoking, asthma, emphysema or if cough is accompanied by excessive phlegm (mucus) unless directed by a physician. Do not exceed recommended dosage. Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. **KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN.** In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Drug Interaction Precaution: Do not use this product if you are now taking a prescription monoamine oxidase inhibitor (MAOI), (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you are uncertain whether your prescription drug contains an MAOI, consult a health professional before taking this product. Store below 86° F and protect from moisture. USE ONLY IF WRAPPER IS UNOPENED.

AVAILABILITY
Sticks of 9 Cough-X® Lozenges.

[HOME](#)[DISCLAIMER / PRIVACY POLICY](#)