

KEYWORD SEARCH

GO

OUR PRODUCTS

STORE LOCATOR

ABOUT US

FAQs

LINKS

CONTACT US



B.F. ASCHER & CO. INC.
pharmaceuticals

*Makers of Consumer Health Care
Products for Better Living.*



Congestac Caplets

- Relieves nasal, sinus, and chest congestion
- Helps clear air passages, making breathing easier
- Does not cause drowsiness

LOCATE THIS PRODUCT IN A STORE NEAR YOU

Click to return to the **PRODUCT CATEGORY PAGE**

Relieves nasal, sinus, and chest congestion without drowsiness

Treat head and chest congestion with Congestac. In convenient caplet form, Congestac contains both a decongestant and expectorant to relieve congestion caused by sinusitis, hay fever, allergies or the common cold. Congestac's effective non-drowsy formula loosens and thins mucus to help drain bronchial tubes and make coughs more productive and ease breathing.

INGREDIENTS

Active Ingredients: Each caplet contains Guaifenesin 400 mg, Pseudoephedrine Hydrochloride 60 mg.
Inactive Ingredients: Croscarmellose Sodium, Hydroxypropyl Methylcellulose, Magnesium Stearate, Microcrystalline Cellulose, Polyethylene Glycol, Povidone, Silicon Dioxide, Starch, and trace amounts of other ingredients

DIRECTIONS

Adults and children over 12: one caplet every 4 to 6 hours, not to exceed 4 caplets in 24 hours.
Children (6 to under 12 years): one-half the adult dose (break caplet in half), not to exceed 2 caplets in 24 hours. Children under 6 years, use only as directed by physician.

Tamper-evident package features for your protection:

Each caplet is encased in a clear plastic cell with a foil back.

The letter "C" appears on each caplet (see product illustration on front of carton).

Do not use this product if either of these tamper-evident features is missing or broken.

USES

- temporarily relieves nasal congestion due to a cold or hay fever, or associated with sinusitis
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

WARNINGS

Do not exceed recommended dosage.

Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions,

or Parkinson's disease) or for two weeks after stopping the MAOI. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Do not take this product for persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema, or where a cough is accompanied by excessive phlegm (mucus) unless directed by a doctor.

A persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by a fever, consult a doctor.

Ask a doctor before use if you have • heart disease • high blood pressure • thyroid disease • diabetes • difficulty in urinating due to an enlarged prostate gland.

Stop and ask a doctor if • you get nervous, dizzy or sleepless • symptoms do not improve within 7 days or are accompanied by fever.

If pregnant or breast-feeding ask a health professional before use.

KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN. In case of overdose, get medical help or contact a poison control center immediately.

Store at room temperature (59°-86° F).

The Congestac horizontal color bar is a trademark of B.F. Ascher Co., Inc.

AVAILABILITY

Congestac caplets are available in packages of 12 and 24 caplets in blisters.

[HOME](#)

[DISCLAIMER / PRIVACY POLICY](#)